



- [Home](#)
- [Clinical Negligence News](#)
- [Clinical Negligence Team](#)
- [About Medical Claims](#)
- [Making a Medical Claim](#)
- [Funding a Medical Claim](#)
- [Inquests](#)
- [Alternatives to Litigation](#)
- [Our Successes](#)
- [Bristol Heart Children](#)
- Netcare -
  - [Orthopaedic Surgery](#)
  - [Ophthalmic Surgery](#)
- [Devon NHS directory](#)
- [Clinical Negligence Links](#)
- [Clinical Negligence FAQ's](#)
- [Contact](#)
- [Michelmores Corporate Website](#)



## Michelmores Medical Negligence News

### vCJD victim had different gene

In: [General](#) On: 18/12/2009

Michelmores' clinical negligence team provide a link to the BBC News online story - click [here](#)

A 30-year-old man thought to have died in January from vCJD belonged to a genetic group that had not shown any signs of the disease, scientists say.

In the UK, 166 people have died of variant CJD, linked to eating BSE-infected beef, and all were thought to have shared a certain gene.

Writing in the Lancet, the scientists say Grant Goodwin, of Lanarkshire, had a different version of the gene.

They estimate that up to 350 people in this group could get vCJD.

Scientists have always thought that a second wave of vCJD cases would emerge some time after the first.

This is the first indication that this theory is being born out with the identification of the first probable vCJD patient outside of the initial genetic group, BBC science correspondent Pallab Ghosh reports.

It is probable because the diagnosis is based on observations of the progression of the disease rather than post-mortem tests which would have provided absolute confirmation of the disease, he adds.

The case report written by Professor John Collinge, of the National Prion Clinic, and colleagues is a reminder that the disease has not gone away.

Many thousands of people may be carrying the infection and although they will never show any symptoms, they have the potential to infect others.

vCJD is caused by infectious agents called prions.

Prion diseases affect the structure of the brain or other neural tissue and are currently untreatable.

Disease-causing prions are thought to consist of abnormally folded proteins, which spread by encouraging the normal healthy prion protein found on the surface of most cells in the body to change shape.

Tests showed that Mr Goodwin had a heterozygous version of the gene which codes for the human prion amino acids valine (V) or methionine (M).

People can be V V (homozygous), M M (homozygous) or M V (heterozygous).

Since 1994, around 200 cases of vCJD have been identified worldwide, and all those tested have been M M homozygous.

However, Mr Goodwin was M V heterozygous.

It is thought that 47% of the population have this version of the gene.

### Navigation

- [Main News](#)
- [Search](#)

### Categories

- [All categories](#)
- [News Archive](#)
- [Our Successes](#)
- [Childbirth, children's injuries & Cerebral Palsy](#)
- [Eye surgery & ophthalmology](#)
- [NHS issues / Outsourcing](#)
- [Heart surgery](#)
- [Cosmetic surgery](#)
- [Meningitis](#)
- [Blood Products Litigation](#)
- [Cancer](#)
- [Product Liability](#)
- [Lawyer Negligence](#)
- [General](#)
- [Spinal Surgery](#)
- [Inquests](#)
- [Brain Injury](#)
- [Overseas Treatment](#)
- [Personal Injury](#)
- [Sports injuries](#)

### Archive

- [December 2009](#)
- [November 2009](#)
- [October 2009](#)
- [September 2009](#)
- [August 2009](#)
- [July 2009](#)
- [June 2009](#)
- [May 2009](#)
- [April 2009](#)
- [March 2009](#)
- [February 2009](#)
- [January 2009](#)
- [December 2008](#)
- [November 2008](#)
- [October 2008](#)
- [September 2008](#)
- [August 2008](#)
- [July 2008](#)
- [June 2008](#)
- [May 2008](#)
- [April 2008](#)
- [March 2008](#)

Professor Collinge said: "The majority of the UK population have potentially been exposed to BSE prions but the extent of clinically silent infection remains unclear.

"About a third of the UK population are M M homozygous.

"If individuals with other genotypes are similarly susceptible to developing prion disease after BSE prion exposure, but with longer incubation periods, further cases would be expected."

The scientists have previously looked at another prion disease in New Guinea, called kuru, which is induced by eating infected human tissues.

The original cases were all M M but more recently M V have appeared.

They say this indicates that M V people can get prion diseases like kuru but have a much longer incubation period.

A Department of Health spokesperson said: "'The Spongiform Encephalopathy Advisory Committee (SEAC) have noted this finding, which confirms the need for ongoing vigilance and robust surveillance of CJD.

"We are continuing to provide resources for CJD surveillance and research, and the development of a test for vCJD remains a priority."

Back to Top  
Tel: 01392 688688  
copyright © Michelmores Solicitors 2003

- February 2008
- January 2008
- December 2007
- November 2007
- October 2007
- September 2007
- August 2007
- July 2007
- June 2007
- May 2007
- April 2007
- March 2007
- February 2007
- January 2007
- December 2006
- November 2006
- October 2006
- September 2006
- August 2006
- July 2006
- June 2006
- May 2006
- April 2006
- March 2006
- February 2006
- January 2006
- December 2005
- November 2005

**Talk to Us Now**

**0800  
0730140**

**Request a call back**

Name:

Phone number:

Email:

Summary of enquiry:

**Send**

**Email Us Now**

Print this page  
Forward to a friend, colleague or adviser

What can you do?

How can we help?